

Veröffentlichungen C. Oberschmidt (geb. Seifried)

Kalbe E, Rehberg SP, Heber I, Kronenbürger M, Schulz JB, Storch A, Linse K, Schneider C, Gräber G, Liepelt-Scarfone I, Berg D, Dams J, Balzer-Geldsetzer M, Hilker R, **Oberschmidt C**, Witt K, Schmidt N, Mollenhauer B, Trenkwalder C, Spoke A, Roeske S, Wittchen HU, Riedel O, Dodel R. Subtypes of mild cognitive impairment in patients with Parkinson's disease: evidence from the LANDSCAPE study. *JNNP*, published on July 8, 2016 as 10.1136/jnnp-2016-313838.

Rodríguez-Labrada R, Velázquez-Pérez L, Aguilera-Rodríguez R, **Seifried-Oberschmidt C**, Peña-Acosta A, Canales-Ochoa N, Medrano-Montero J, Estupiñan-Rodríguez A, Vázquez-Mojena Y, González-Zaldivar Y, Laffita Mesa JM. Executive deficit in spinocerebellar ataxia type 2 is related to expanded CAG repeats: Evidence from antisaccadic eye movements. *Brain Cogn*. 2014 Nov;91:28-34.

Baudrexel S, **Seifried C**, Penndorf B, Klein JC, Middendorp M, Steinmetz H, Grünwald F, Hilker R. The value of putaminal diffusion imaging versus 18-fluorodeoxyglucose positron emission tomography for the differential diagnosis of the Parkinson variant of multiple system atrophy. *Mov Disord*. 2014 Mar;29(3):380-7.

Arnold C, Gehrig J, Gispert S, **Seifried C**, Kell CA. Pathomechanisms and compensatory efforts related to Parkinsonian speech. *Neuroimage Clin*. 2013 Oct 31;4:82-97.

Weise LM, **Seifried C**, Eibach S, Gasser T, Roeper J, Seifert V, Hilker R. Correlation of active contact positions with the electrophysiological and anatomical subdivisions of the subthalamic nucleus in deep brain stimulation. *Stereotact Funct Neurosurg*. 2013;91(5):298-305.

Seifried C, Boehncke S, Heinzmann J, Baudrexel S, Weise L, Gasser T, Eggert K, Fogel W, Baas H, Badenhoop K, Steinmetz H, Hilker R. Diurnal variation of hypothalamic function and chronic subthalamic nucleus stimulation in Parkinson's disease. *Neuroendocrinology*. 2013;97(3):283-90.

Damrath E, Heck MV, Gispert S, Azizov M, Nowock J, **Seifried C**, Rüb U, Walter M, Auburger G. ATXN2-CAG42 sequesters PABPC1 into insolubility and induces FBXW8 in cerebellum of old ataxic knock-in mice. *PLoS Genet*. 2012;8(8):e1002920.

Seifried C, Weise L, Hartmann R, Gasser T, Baudrexel S, Szelényi A, van de Loo S, Steinmetz H, Seifert V, Roeper J, Hilker R. Intraoperative microelectrode recording for the delineation of subthalamic nucleus topography in Parkinson's disease. *Brain Stimul*. 2012 Jul;5(3):378-87.

Klein JC, Barbe MT, **Seifried C**, Baudrexel S, Runge M, Maarouf M, Gasser T, Hattingen E, Liebig T, Deichmann R, Timmermann L, Weise L, Hilker R. The tremor network targeted by successful VIM deep brain stimulation in humans. *Neurology*. 2012 Mar 13;78(11):787-95.

Roggendorf J, Chen S, Baudrexel S, van de Loo S, **Seifried C**, Hilker R. Arm swing asymmetry in Parkinson's disease measured with ultrasound based motion analysis during treadmill gait. *Gait Posture*. 2012 Jan;35(1):116-20.

Oberschmidt C, Hilker R, Kang JS. Neues zu Neurologischen Bewegungsstörungen 2011. *Akt Neurologie* 2011 ; 38:222-233.

Rodríguez-Labrada R, Velázquez-Pérez L, **Seifried C**, Canales-Ochoa N, Auburger G, Medrano-Montero J, Sánchez-Cruz G, Aguilera-Rodríguez R, Lafitta-Mesa J, Vázquez-Mojena Y, Verdecia-Ramirez M, Motta M, Quevedo-Batista Y. Saccadic latency is prolonged in Spinocerebellar Ataxia type 2 and correlates with the frontal-executive dysfunctions. *J Neurol Sci* 2011;306:103-107.

Balzer-Geldsetzer M, Ferreira Braga da Costa AS, Kronenbürger M, Schulz JB, Röske S, Wüllner U, Klockgether T, Storch A, Schneider C, Riedel O, Wittchen HU, **Seifried C**, Hilker R, Schmidt N, Witt K, Deuschl G, Mollenhauer B, Trenkwalder C, Liepelt-Scarfone I, Berg D, Gasser T, Kalbe E, Bodden M, Oertel WH, Dodel R. Parkinson's disease dementia: a longitudinal study (DEMPARK). *Neuroepidemiology* 2011;37(3-4):168-76.

Baudrexel S, Witte T, **Seifried C**, von Wegner F, Beissner F, Klein JC, Steinmetz H, Deichmann R, Roeper J, Hilker R. Resting state fMRI reveals increased subthalamic nucleus-motor cortex connectivity in Parkinson's disease. *Neuroimage*. 2011 Apr 15;55(4):1728-38.

Klein JC, Lorenz B, Kang JS, Baudrexel S, **Seifried C**, van de Loo S, Steinmetz H, Deichmann R, Hilker R. Diffusion tensor imaging of white matter involvement in essential tremor. *Hum Brain Mapp.* 2011 Jun;32(6):896-904.

Hattingen E, Magerkurth J, Pilatus U, Mozer A, **Seifried C**, Steinmetz H, Zanella F, Hilker R. Phosphorus and proton magnetic resonance spectroscopy demonstrates mitochondrial dysfunction in early and advanced Parkinson's disease. *Brain.* 2009 Dec;132(Pt 12):3285-97.

Velázquez-Pérez L, **Seifried C**, Abele M, Wirjatijasa F, Rodríguez-Labrada R, Santos-Falcón N, Sánchez-Cruz G, Almaguer-Mederos L, Tejeda R, Canales-Ochoa N, Fetter M, Ziemann U, Klockgether T, Medrano-Montero J, Rodríguez-Díaz J, Laffita-Mesa JM, Auburger G. Saccade velocity is reduced in presymptomatic Spinocerebellar Ataxia Type 2, *Clin Neurophysiol.* 2009 Mar;120(3):632-5.

Fernandez-Ruiz J, Velázquez-Perez L, Díaz R, Drucker-Colín R, Pérez-González R, Canales N, Sánchez-Cruz G, Martínez-Góngora E, Medrano Y, Almaguer-Mederos L, **Seifried C**, Auburger G. Prism adaptation in spinocerebellar ataxia type 2. *Neuropsychologia.* 2007 Sep 20;45(12):2692-8.

Seifried C, Weidauer S, Hinsch N, Steinmetz H. Cerebrale lymphomatoid granulomatosis: A case report. *Nervenarzt.* 2007 Feb;78(2):193-7.

Otto B, Hermans M, **Seifried C**, Buchkremer M, Lanfermann H, Sitzer M. Neurosyphilis Important differential diagnosis of herpes simplex encephalitis. *Nervenarzt.* 2005 Oct 14; 2007 Aug;78(8):944-7.

Seifried C, Sitzer M, Jen J, Auburger G. HERNs: a rare, hereditary, multisystemic disease with cerebral microangiopathy. *Nervenarzt.* 2005 Oct;76(10):1191-2, 1194-5.

Seifried C, Velázquez-Pérez L, Santos-Falcón N, Abele M, Ziemann U, Almaguer LE, Martínez-Góngora E, Sánchez-Cruz G, Canales N, Pérez-González R, Velázquez-Manresa M, Viebahn B, Stuckrad-Barre S, Klockgether T, Fetter M, Auburger G. Saccade velocity as a surrogate disease marker in spinocerebellar ataxia type 2. *Ann NY Acad Sci.* 2005 Apr;1039:524-7.

Velázquez-Pérez L, **Seifried C**, Santos-Falcón N, Abele M, Ziemann U, Almaguer LE, Martínez-Góngora E, Sánchez-Cruz G, Canales N, Pérez-González R, Velázquez-Manresa M, Viebahn B, Stuckrad-Barre S, Fetter M, Klockgether T, Auburger G. Saccade velocity is strongly influenced by polyglutamine size in spinocerebellar ataxia 2. *Ann Neurol.* 2004 Sep;56(3):444-7.

Seifried C, Weber M, Bialleck H, Seifried E, Schrezenmeier H, Roth WK. High prevalence of GBV-C/HGV among relatives of GBV-C/HGV-positive blood donors in blood recipients and in patients with aplastic anemia Transfusion. 2004 Feb;44(2):268-74.

Rüb U, Del Turco D, Del Tredici K, de Vos RA, Brunt ER, Reifenberger G, **Seifried C**, Schultz C, Auburger G, Braak H. Thalamic involvement in a spinocerebellar ataxia type 2 (SCA2) and a spinocerebellar ataxia type 3 (SCA3) patient, and its clinical relevance. *Brain.* 2003 Oct;126(Pt 10):2257-72.

Rüb U, Schultz C, Del Tredici K, Gierga K, Reifenberger G, de Vos RA, Braak H, Auburger G. Rub U, Schultz C, Del Tredici K, Gierga K, Reifenberger G, de Vos RA, **Seifried C**, Braak H, Auburger G. Thalamic involvement in a spinocerebellar ataxia type 2 (SCA2) and a spinocerebellar ataxia type 3 (SCA3) patient, and its clinical relevance *Neuropathol Appl Neurobiol.* 2003 Oct;29(5):418-33.